arXiv:1106.0348v1 [math.RA] 2 Jun 2011

On zero divisors and prime elements of po-semirings

Tongsuo Wu[a]*, Dancheng Lu[b]†, Yuanlin Li[c]‡

[a]Department of Mathematics, Shanghai Jiaotong University
Shanghai 200240, P. R. China

[b]Department of Mathematics, Suzhou University, Suzhou 215006, P.R. China

[c]Department of Mathematics, Brock University, St. Catharines, On., Canada L2S 3A1

Abstract. A semiring is an algebraic structure similar to a ring, but without the requirement that each element must have an additive inverse. A po-semiring is a semiring equipped with a compatible bounded partial order. In this paper, properties of zero divisors and prime elements of a po-semiring are studied. In particular, it is proved that under some mild assumption the set $Z(A)$ of nonzero zero divisors of A is $A\setminus\{0,1\}$, each prime element of A is a maximal element, and the zero divisor graph $\Gamma(A)$ of A is a finite graph if and only if A is finite. For a po-semiring A with $Z(A)=A\setminus\{0,1\}$, it is proved that A has finitely many maximal elements if ACC holds either for elements of A or for principal annihilating ideals of A. As applications of prime elements, it is shown that the structure of a po-semiring A is completely determined by the structure of integral po-semirings if either $|Z(A)|=1$ or $|Z(A)|=2$ and $Z(A)^2\neq 0$. Applications to the ideal structure of commutative rings are considered.

Key Words: po-semiring, zero divisor, prime element, small $Z(A)$, graph property

1. Introduction and preliminaries

Throughout this paper, all semigroups S and all rings R are assumed to be commutative with zero element 0_S (i.e., $0S=0$) and with identity 1_R respectively. For a semigroup S, let $Z(S)$ be the set of nonzero zero divisors and S^* the set of nonzero elements of S. For

---

[*] Corresponding author, tswu@sjtu.edu.cn (T. Wu)

[†] ludancheng@suda.edu.cn (D. Lu)

[‡] yli@brocku.ca (Y. Li)

a ring R, let $U(R)$ be the set of invertible elements of R, $N(R)$ the nil radical and $J(R)$ the Jacobson radical of R.

A *commutative semiring* is a set A which contains at least two elements $0, 1$ and which is equipped with two binary operations, $+$ and $\cdot$, called addition and multiplication respectively, such that the following conditions hold:

1. $(A, +, 0)$ is a commutative monoid with zero element 0.
2. $(A, \cdot, 1)$ is a commutative monoid with identity element 1.
3. Multiplication distributes over addition.
4. 0 annihilates A with respect to multiplication, i.e., $0a = 0$, $\forall a \in A$.

If there is no nonzero zero-divisor in a semiring A, then A is called an *integral semiring*. Certainly, each ring is a semiring. Other important examples of semirings include the set $\mathbb{I}(R)$ of ideals of a commutative ring R, the set $\mathbb{N}$ of nonnegative integers, and the real segment $[0, 1]$ whose addition is *max* operation. Note that both $\mathbb{N}$ and $[0, 1]$ are integral semirings. Based on a semiring structure, an excellent and rather general framework for constraint satisfaction and optimization was developed in [6, 7].

Next we introduce a new notion which will be the central topic of this paper.

Definition 1.1. A *partially-ordered semiring* is a commutative semiring $(A, +, \cdot, 0, 1)$, together with a compatible partial order $\leq$, i.e., a partial order $\leq$ on the underlying set A that is compatible with the semiring operations in the sense that it satisfies the following conditions:

(1) $x \leq y$ implies $x + z \leq y + z$, and

(2) $0 \leq x$ and $y \leq z$ imply that $xy \leq xz$ for all x, y, z in A.

If A satisfies the following additional condition, then A is called a *po-semiring*:

(3) The partial set $(A, \leq, 0, 1)$ is bounded, i.e., 1 is the largest element and 0 is the least element of A.

We remark that condition (3) is so strong that it enables a po-semiring A to be a dioid, where a semiring is called a *dioid* if its addition is idempotent ($a + a = a$, $\forall a \in A$). Furthermore, the above defined partial order $\leq$ for a po-semiring A is identical with the new partial order $\leq_1$ defined by the following

$$a \leq_1 b \text{ if and only if } a + b = b.$$

In other words, $(A, +, 0, 1)$ is a bounded join-semilattice. Clearly, any bounded, distributive lattice is a commutative dioid under join and meet, where $a \leq b$ iff $a \wedge b = a$. Each bounded, distributive lattice is certainly also a po-semiring under the Definition 1.1.

An element p of a po-semiring is called *prime*, if $p \neq 1$ and $xy \leq p$ implies either $x \leq p$ or $y \leq p$. An element x is called *minimal*, if $x \neq 0$ and $0 < y \leq x$ implies $x = y$. An element $\mathfrak{m}$ is called *maximal*, if $\mathfrak{m} \neq 1$ and $\mathfrak{m} \leq x < 1$ implies $\mathfrak{m} = x$.

An *ideal* I of a po-semiring A is an additive sub-semigroup containing the zero element

0_A such that $IA \subseteq I$. For each element u of a po-semiring A, set

$$<u> = \{x \in A \,|\, x \leq u\}.$$

Clearly, $<u>$ is an ideal of A, called the *lower principle ideal generated by* u. The principle ideal uA is certainly another ideal generated by u, and clearly $uA \subseteq <u>$. An ideal I of A is called *hereditary*, if $<u> \subseteq I$ holds for all u in I. For any element u of A, both $<u>$ and $ann_A(u)$ are hereditary ideals of A.

Our prototype of po-semiring is the po-semiring $\mathbb{I}(R)$ of a commutative ring R, which consists of all ideals of R. The multiplication is the ideal multiplication, and the addition is the addition of subsets. The partial order is the usual inclusion. The po-semiring $\mathbb{I}(R)$ also has the property that for each element $v \neq 1$, there exists a maximal element $\mathfrak{m}$ such that $v \leq \mathfrak{m}$. In $\mathbb{I}(R)$, there is the notion of infinite sum of elements of $\mathbb{I}(R)$. Notice that $\mathbb{I}(R)$ is also a bounded lattice under $I \wedge J = I \cap J$ and $I \vee J = I + J$, but it is not necessarily a distributive lattice for a general ring R. Denote $\{0, 1\} = \mathbb{I}(F)$ for any field F, and note that it is the smallest po-semiring, i.e., it can be embedded into any po-semiring.

Throughout the paper, we always assume that one of the following three additional conditions on a po-semiring A holds:

(C_1): *For each non-nilpotent element u of A, there is a nonzero idempotent w such that $w \leq u$ and w has an orthogonal idempotent complement in A, i.e., there exists an idempotent element v in A such that $w + v = 1, wv = 0$.*

(C_2): *For each nonzero idempotent element u of A, there is a nonzero idempotent w such that $w \leq u$ and w has an orthogonal idempotent complement in A.*

(C_3): *Each idempotent minimal element of A has an orthogonal idempotent complement in A.*

Clearly, $(C_1) \Longrightarrow (C_2) \Longrightarrow (C_3)$ hold for any po-semiring A. In sections 2 and 3, examples will be given to show that the reverse implications do not hold. Note that if $Z(A) \neq \emptyset$, then each minimal element of A is a zero divisor (see Lemma 3.4(2)).

Proposition 1.2. *For any commutative ring R, let $A = \mathbb{I}(R)$. Then A satisfies the condition (C_3).*

(1) *If further R is a noetherian exchange ring, then A satisfies condition (C_2).*

(2) *If further R is a noetherian exchange ring and $J(R) = N(R)$, then A satisfies condition (C_1). In particular, it holds for any artinian ring R.*

(3) *If further A satisfies condition (C_1), then $J(R) = N(R)$.*

Proof. The first conclusion follows from Brauer's Lemma, see [9, 10.22].

(1) Let I be a nontrivial idempotent ideal of R. Then I is finitely generated, and thus $I \not\subseteq J(R)$ by Nakayama Lemma. Then the exchange property ensures the existence of a

nonzero idempotent e in I. Thus $(Re)^2 = Re$, Re is a nonzero idempotent and it has the orthogonal idempotent complement $R(1-e)$. Hence (C_2) holds for $\mathbb{I}(R)$.

(2) An artinian ring R is a noetherian exchange ring and each non-nilpotent ideal of R contains a nonzero idempotent.

(3) For any $x \notin N(R)$, Rx is not nilpotent. By condition (C_1), there exists a nonzero idempotent e in Rx. Thus $x \notin J(R)$. This shows $J(R) = N(R)$. $\blacksquare$

We remark that exchange rings include von Neumann regular rings, artinian rings, semilocal rings such that idempotents lift modulo their Jacobson radical. For further information on exchange rings, see [10] and the listed references.

In this paper we investigate properties of zero divisors and prime elements of a po-semiring. Section 2 deals with the following questions: (1) Under what conditions can it occur that $Z(A) = A \setminus \{0,1\}$? (2) When does a po-semiring A have only finitely many maximal elements? (3) When is every prime element also maximal? It is proved that $Z(A) = A \setminus \{0,1\}$, each prime element of A is a maximal element, if one of the following conditions is satisfied: (i) Condition (C_2) holds in A, and DCC holds for elements of A. (ii) Condition (C_1) holds in A, and there exists no infinite set of orthogonal idempotents in A. For a po-semiring A with $Z(A) = A \setminus \{0,1\}$, it is also proved that A has finitely many maximal elements if ACC holds either for elements of A or for principal annihilating ideals of A. In Section 3, we study the zero divisor graph $\Gamma(A)$ of a po-semiring A. In particular, for a po-semiring A satisfying condition (C_3), it is proved that the graph $\Gamma(A)$ is either a star graph or a two-star graph $K_1 + K_1 + K_1 + D_r$ if G contains no cycle. It is proved that $\Gamma(A) \cong K_1 + K_1 + K_1 + D_r$ if and only if $A \cong \{0,1\} \times S$, where S is a po-semiring with $|Z(S)| = 1$ (in this case, $r = |S| - 2$). Example 3.6 shows further that for any r with $1 \leq r \leq \aleph_0$, there exists a po-semiring A such that condition (C_3) holds and $\Gamma(A) \cong K_1 + K_1 + K_1 + D_r$. In Section 4, we study the structure of a po-semiring A with $1 \leq |Z(A)| \leq 2$. It is shown that the structure of a po-semiring A is completely determined by the structures of integral po-semirings, if either $|Z(A)| = 1$ or $|Z(A)| = 2$ and $Z(A)^2 \neq 0$ (Theorem 4.2). When A is taken to be $\mathbb{I}(R)$ for some commutative ring R, applications to the ideal structure of a ring are provided.

2. Chain conditions on po-semirings

In this section, we study properties of elements of $A \setminus \{0,1\}$. We begin with an easy proposition which will be used repeatedly.

Proposition 2.1. *Let A be a po-semiring. Then*

(1) *Each maximal element of A is prime.*

(2) *For a maximal element $\mathfrak{m}$ of A, if $\mathfrak{m}b = 0$, then either $\mathfrak{m}^2 = \mathfrak{m}$ or $b^2 = 0$.*

(3) *If* $e_i + f_i = 1$ *where* $e_if_i = 0$, $e_i^2 = e_i$, $f_i^2 = f_i$, *then* $e_1 > e_2$ *implies* $f_1 < f_2$.

Proof. (1) Let $\mathfrak{m}$ be a maximal element of A. Assume that $ab \leq \mathfrak{m}$ and $a \not\leq \mathfrak{m}$. Then $a + \mathfrak{m} > \mathfrak{m}$, so $a + \mathfrak{m} = 1$. Then $b = ab + \mathfrak{m}b \leq \mathfrak{m} + \mathfrak{m} = \mathfrak{m}$, so $\mathfrak{m}$ is a prime element of A.

(2) For a maximal element $\mathfrak{m}$ of A and any $b \in A$, either $\mathfrak{m} + b = \mathfrak{m}$ or $\mathfrak{m} + b = 1$. If $\mathfrak{m} + b = \mathfrak{m}$, then $b^2 = 0$. If $\mathfrak{m} + b = 1$, then $\mathfrak{m} = \mathfrak{m}^2$.

(3) If $e_1 \geq e_2$, then $e_1 + f_2 \geq e_2 + f_2 = 1$, and hence $f_1 \leq f_2f_1 \leq f_2$. If $f_1 = f_2$, then $e_1 = e_1(e_2 + f_2) = e_1e_2 \leq e_2$. This completes the verification. ■

Recall that a nonzero idempotent of a semiring is called a *primitive idempotent* if it cannot be written as a sum of two orthogonal nontrivial idempotents.

Theorem 2.2. *For a po-semiring* A, $Z(A) = A \setminus \{0, 1\}$ *if one of the following conditions holds:*

(1) *Condition* (C_1) *holds and there exists no infinite set of orthogonal idempotents in* A.

(2) *DCC holds for elements of* A, *and condition* (C_2) *holds.*

In both cases, for any element c *in* $A \setminus \{0, 1\}$, *either* c *is nilpotent or there exist a positive integer* n *and a nontrivial idempotent* e *such that* $c^n = c^ne$, *where* e *has an orthogonal idempotent complement in* A.

Proof. (1) Suppose that condition (C_1) holds in A. Assume to the contrary that there exists an element a in $A \setminus \{0, 1\}$ such that $a \notin Z(A)$. Then a is not nilpotent. Hence by condition (C_1), there exist nonzero idempotents e_1, f_1 such that $e_1 \leq a, e_1f_1 = 0, e_1 + f_1 = 1$. If af_1 is nilpotent, then a is a zero divisor. If further a is idempotent, then $af_1 = 0$ and thus $a = ae_1 \leq e_1$, so $a = e_1$.

Note that af_1 is not nilpotent by assumption on a. Again by condition (C_1), there exist nonzero orthogonal idempotents e_2, f_2 such that $e_2 + f_2 = 1, e_2 \leq af_1$. Then $e_1e_2 = 0$, so $e_1 < e_1 + e_2$ and $(e_1 + e_2) + (f_1f_2) = 1$. Note that f_1f_2 is a nonzero idempotent and is orthogonal to the idempotent $(e_1 + e_2)$. If $a(f_1f_2)$ is nilpotent, then a is a zero divisor. If further a is idempotent, then $af_1f_2 = 0$, implying $a = a(e_1 + e_2) \leq e_1 + e_2$, so $a = e_1 + e_2$.

Clearly, $a(f_1f_2)$ is not nilpotent by assumption on a. Then there exist nonzero orthogonal idempotents e_3, f_3 such that $e_3 + f_3 = 1, e_3 \leq a(f_1f_2)$. Then we have an orthogonal idempotent decomposition $(e_1+e_2+e_3)+(f_1f_2f_3) = 1$, and thus $e_1 < e_1+e_2 < e_1+e_2+e_3$, where $e_ie_j = \delta_{ij}e_i$. Finally, if a is idempotent and $a(f_1f_2f_3) = 0$, then $a = e_1 + e_2 + e_3$. Clearly, $a(f_1f_2f_3)$ is not nilpotent under the assumption on a.

Continuing this process, we finally obtain an infinite set $\{e_1, e_2, \cdots, \}$ of orthogonal idempotents in A, contradicting the assumption on A. Note that $e_1 < e_1 + e_2 < e_1 + e_2 + e_3 < \cdots$ and $(e_1 + \cdots + e_i) + (f_1 \cdots f_i) = 1$ implies $f_1 > f_1f_2 > \cdots$ by Lemma 2.1(3). Note also that $e_1 + \cdots + e_i \leq a$. Note that if assume $a(f_1f_2 \cdots f_i) = 0$ for some i, then $a = e_1 + e_2 + \cdots + e_i$.

(2) Assume the condition (C_2) holds for A and DCC holds for elements of A. Assume to the contrary that in $A \setminus \{0, 1\}$ there exists an element b such that $b \notin Z(A)$. Then b is not nilpotent and there exists a positive integer m such that b^m is idempotent. Let $a = b^m$. By repeating the discussions in the proof in part (1), we obtain an infinite descending chain of elements $f_1 > f_1 f_2 > \cdots$, giving a contradiction. ■

By the proof of Theorem 2.2, we have the following improved result for idempotent elements.

Theorem 2.3. *For a po-semiring A, if condition (C_2) holds in A and A contains no infinite set of orthogonal idempotents, then each idempotent element of A has an orthogonal idempotent complement. In particular, each nontrivial idempotent is a zero divisor of A. Furthermore, each nonzero idempotent of A is a finite sum of orthogonal primitive idempotents.*

Proof. The result follows from the proof of (1) in Theorem 2.2 if we start with a nontrivial idempotent a. ■

We remark that there exists no infinite set of orthogonal idempotents in a po-semiring A provided that one of the following conditions holds: (1) ACC holds for idempotent elements of A. (2) DCC holds for idempotent elements of A, and condition (C_2) also holds. The latter conclusion follows from Theorem 2.3 and Lemma 2.1(3). If ACC (respectively, DCC) holds for elements of A, then for each element $a \neq 0, 1$, clearly there is a maximal (respectively, minimal) element x of A such that $x \geq a$ (respectively, $x \leq a$).

Let $A = \mathbb{I}(R)$ for some commutative ring R. In view of Theorems 2.2 and 2.3, we have the following applications to commutative rings.

Corollary 2.4. *Assume that a commutative ring R satisfies one of the following conditions:*

(1) *Each non-nilpotent ideal of R contains a nonzero idempotent, and R contains no infinite set of orthogonal idempotents.*

(2) *R is artinian.*

Then for each nontrivial ideal I of R, there exists a nontrivial idempotent element e such that $I \subseteq Re$. Furthermore, each nonzero idempotent ideal of R has the form $\sum_{i=1}^{r} Re_i$, where $e_1, \cdots, e_r$ are orthogonal primitive idempotents.

Corollary 2.5. *For a commutative noetherian exchange ring R, each nontrivial idempotent ideal of R is an annihilating ideal. Furthermore, each nonzero idempotent ideal of R has the form $\sum_{i=1}^{r} Re_i$, where $e_1, \cdots, e_r$ are orthogonal primitive nonzero idempotents.*

Proof. By applying Proposition 1.2(1) and Theorem 2.3 to $\mathbb{I}(R)$, we obtain the result.

■

Note that any artinian ring is a noetherian exchange ring. Thus Corollary 2.5 holds in particular for artinian rings.

It is well-known that in a commutative artinian ring R, each prime ideal is a maximal ideal of R, i.e., each prime element of the po-semiring $\mathbb{I}(R)$ is a maximal element if DCC holds for elements of $\mathbb{I}(R)$. Also, if DCC holds for elements of $\mathbb{I}(R)$, then ACC also holds. The following example shows that the above mentioned results are not true for a general po-semiring. It also shows that the additional condition (C_2) in Theorem 2.3 is needed, and that $Z(A) = A \setminus \{0, 1\}$ does not imply condition (C_2).

Example 2.6 *There exists an infinite po-semiring A such that DCC holds but ACC fails for elements of A, $Z(A) = A \setminus \{0, 1\}$ and A has infinitely many prime elements but none of which is a maximal element.*

Let $A = \{0, 1, a, b_1, b_2, \cdots\}$ be a countable set with $4 \leq |A| \leq \aleph_0$, and define a partial order $\leq$ by $0 < a < b_1 < b_2 < \cdots < 1$ on A. Define an addition by $x + y = \max\{x, y\}$, $\forall x, y \in A$. Define a commutative multiplication by

$$0x = 0,\ 1x = x\ (\forall x \in A),\ ab_i = 0,\ a^2 = 0,\ b_i b_j = b_{min\{i,j\}}.$$

Then it is routine to check that A is a po-semiring. Note that a and b_i are prime elements of A for all i, and condition (C_2) does not hold for A although condition (C_3) does hold. The elements of A satisfies DCC, but they do not satisfy ACC if $|A|$ is infinite. Note also that $Z(A) = A \setminus \{0, 1\}$, and there exists no infinite set of orthogonal idempotents.

On the other hand, for a noetherian ring R which is not artinian, ACC holds for elements of the po-semiring $\mathbb{I}(R)$ but DCC fails.

Despite of the above example, we are able to show that under some suitable condition the set of the maximal elements is the same as the set of the prime elements of A.

Theorem 2.7 *For a po-semiring A, each prime element of A is a maximal element if one of the following condition holds:*

(1) *Condition (C_2) holds in A, and DCC holds for elements of A.*

(2) *Condition (C_1) holds in A, and there exists no infinite set of orthogonal idempotents in A.*

Proof. (1) Assume to the contrary that there exists a prime element q such that $q < p$ for some $p < 1$. Then p is not nilpotent. By the DCC assumption, there exists a positive integer r such that p^r is nonzero and idempotent. By Theorem 2.3, p^r is annihilated by a nonzero idempotent, say f.

By condition (C_2), there exist orthogonal nonzero idempotents f_1, g_1 such that $f_1 \leq f$, $g_1 + f_1 = 1$. Then $p^r \cdot f_1 = 0$. Since $q < p$ and q is prime, we have $f_1 \leq q$ and hence $g_1 \not\leq q$. Clearly, $p^r g_1 = p^r \not\leq q$ and it is idempotent.

By condition (C_2), there exist nonzero orthogonal idempotents t_2, s_2 such that $t_2 \leq p^r g_1$, $t_2 + s_2 = 1$. Then we have $f_1 t_2 = 0$. This together with $t_2 + s_2 = 1$ and $f_1 + g_1 = 1$ implies $f_1 + t_2 + g_1 s_2 = 1$, where $f_1, t_2, g_1 s_2$ are mutually orthogonal idempotents. If $s_2 \not\leq q$, then $t_2 \leq q$. In this case, let $f_2 = t_2$, $g_2 = g_1 s_2$. If $s_2 \leq q$, then $t_2 \not\leq q$ and $g_1 s_2 \leq q$. In this case, let $f_2 = g_1 s_2, g_2 = t_2$. In either case, we have an orthogonal idempotent decomposition $f_1 + f_2 + g_2 = 1$, where $f_1 \leq q$, $f_2 \leq q$ and $g_2 \not\leq q$.

The next step is to consider the idempotent $p^r g_1 g_2$. Clearly $p^r g_1 g_2 \not\leq q$ and in particular, $p^r g_1 g_2 \neq 0$. By condition (C_2), there exist orthogonal idempotents t_3, s_3 such that $t_3 + s_3 = 1$ and $t_3 \leq p^r g_1 g_2$. Clearly, $t_3 f_i = 0$ for $i = 1, 2$. This together with $(f_1 + f_2) + g_2 = 1$ implies $f_1 + f_2 + t_3 + s_3 g_2 = 1$. Since $f_1 + f_2 + t_3 \leq p$, $s_3 g_2 \neq 0$. If $s_3 \not\leq q$, then $t_3 \leq q$. In this case, let $f_3 = t_3$, $g_3 = s_3 g_2$. If $s_3 \leq q$, then $t_3 \not\leq q$ and $g_2 s_3 \leq q$. In this case let $f_3 = g_2 s_3$, $g_3 = t_3$. In either case, we have an orthogonal idempotent decomposition $(f_1 + f_2 + f_3) + g_3 = 1$, where $f_i \leq q$ $(\forall i = 1, 2, 3)$ and $g_3 \not\leq q$.

Continuing this process, we have got an infinite set of mutually orthogonal idempotents $\{f_1, f_2, \cdots\}$. Since $f_1 < f_1 + f_2 < f_1 + f_2 + f_3 < \cdots$, by Lemma 2.1(3) we have an infinite descending chain of idempotents $g_1 > g_2 > \cdots$, contradicting the DCC assumption.

(2) Assume to the contrary that there exists a prime element q such that $q < p$ for some $p < 1$. Then p is not nilpotent. By Theorem 2.2, there exists a positive integer r such that p^r is annihilated by a nonzero idempotent, say, f. The rest of the proof is almost the same as in (1). Note that $p^r(g_1 g_2 \cdots g_s) \not\leq q$ implies that $p^r(g_1 g_2 \cdots g_n)$ is not nilpotent and thus condition (C_1) applies. ■

Corollary 2.8. *For a commutative ring R, each prime ideal of R is a maximal ideal if one of the following condition is satisfied:*

(1) *There exists no infinite set of orthogonal idempotents in R and each non-nilpotent ideal contains a nonzero idempotent.*

(2) *R is an exchange ring, $J(R) = N(R)$ and there exists no infinite set of orthogonal idempotents in R.*

(3) *R is an artinian ring.*

Call an ideal I of a po-semiring A a *principal annihilating ideal* if $I = ann_A(u)$ for some element u of A. The proof of the following theorem is a typical example of use of ideas in proving the Chinese Remainder Theorem.

Theorem 2.9. *For a po-semiring A, assume $Z(A) = A \setminus \{0, 1\}$. If ACC holds either for elements of A or for principal annihilating ideals of A, then A has finitely many maximal elements.*

Proof. Suppose that $Z(A) = A \setminus \{0, 1\}$. Assume to the contrary that A has infinitely many maximal elements and let $\mathfrak{m}_i$ $(i \in N^+)$ be distinct maximal elements of A. Then $\mathfrak{m}_1 + \mathfrak{m}_3 = 1$ and $\mathfrak{m}_2 + \mathfrak{m}_3 = 1$ and hence $1 = \mathfrak{m}_3(\mathfrak{m}_1 + \mathfrak{m}_2 + \mathfrak{m}_3) + \mathfrak{m}_1\mathfrak{m}_2 = \mathfrak{m}_3 + \mathfrak{m}_1\mathfrak{m}_2$. In a similar way, we obtain $\mathfrak{m}_1\mathfrak{m}_2\cdots\mathfrak{m}_m+\mathfrak{m}_{m+1}=1$. Now consider the following ascending chain of principal annihilating ideals of the semiring A

$$ann_A(\mathfrak{m}_1)\subseteq\cdots\subseteq ann_A(\mathfrak{m}_1\cdots\mathfrak{m}_m)\subseteq ann_A(\mathfrak{m}_1\cdots\mathfrak{m}_m\mathfrak{m}_{m+1})\subseteq\cdots.$$

If ACC holds for principal annihilating ideals of A, then there exists an integer m such that $ann_A(\mathfrak{m}_1\cdots\mathfrak{m}_m)=ann_A(\mathfrak{m}_1\cdots\mathfrak{m}_m\mathfrak{m}_{m+1})$. By assumption, there is a nonzero element $x\in ann_A(\mathfrak{m}_{m+1})$. But then $x=x\mathfrak{m}_{m+1}+x(\mathfrak{m}_1\cdots\mathfrak{m}_m)=0$, a contradiction.

If ACC holds for elements of A, we claim that there exists no strict ascending chain $ann_A(\mathfrak{m}_1)<ann_A(\mathfrak{m}_1\mathfrak{m}_2)<\cdots$, and the result thus follows. In fact, if this were not the case, then for each m there would exist an element y_{m+1} such that

$$y_{m+1}\in ann_A(\mathfrak{m}_1\cdots\mathfrak{m}_m\mathfrak{m}_{m+1})\setminus ann_A(\mathfrak{m}_1\cdots\mathfrak{m}_m).$$

Then we would have obtained an infinite ascending chain of elements of A:

$$y_2<y_2+y_3<y_2+y_3+y_4<\cdots.$$

This completes the proof. ■

Combining Theorems 2.2, 2.7 and 2.9, we have the following results.

Corollary 2.10. *Let A be a po-semiring satisfying condition (C_2). If both ACC and DCC holds for elements of A, then A has only a finite number of prime elements. These primes are precisely the maximal elements of A. Under the assumption, $Z(A)=A\backslash\{0,1\}$.*

Corollary 2.11. *Let A be a po-semiring satisfying condition (C_1). If ACC holds for elements of A, then A has only a finite number of maximal elements.*

We remark that if a po-semiring A has a unique maximal element $\mathfrak{m}$, then $\mathfrak{m}$ is nilpotent if condition (C_1) is further assumed to hold in A.

Applying Theorems 2.9 to $\mathbb{I}(R)$ for a commutative ring R, we have the following.

Corollary 2.12. *Let R be a commutative noetherian ring.*

(1) ([5, [Proposition 1.7]) *If each nontrivial ideal is an annihilating ideal, then R is a semilocal ring.*

(2) *If each non-nilpotent ideal of R contains an idempotent, then R is a semilocal ring.*

(3) *Any Noetherian exchange ring R with $J(R)=N(R)$ is a semilocal ring.*

Recall that an ideal I of a po-semiring A is called *hereditary*, if $<u>\subseteq I$ holds for all u in I. Motivated by Theorem 2.9 we have the following.

Proposition 2.13 *Let A be a po-semiring. Then*

(1) *ACC holds for elements of A if and only if ACC holds for hereditary ideals of A.*

(2) *If DCC holds for hereditary ideals of A, then DCC also holds for elements of A. The converse holds, if there is the concept of an infinite sum of elements in A and any hereditary ideal I is closed under taking infinite sums.*

Proof. $\Longleftarrow$ of (1) and (2): For any $u \in A$, $< u >$ is clearly a hereditary ideal of the po-semiring A. For elements u, v of A, $u \leq v$ if and only if $< u > \subseteq < v >$, and $u < v$ if and only if $< u > \subset < v >$. This implies the sufficiency part of the proposition.

$\Longrightarrow$: (1) If ACC does not hold for hereditary ideals of the po-semiring A, then there exists a strict ascending chain of hereditary ideals of A, say, $X_1 \subset X_2 \subset \cdots$. Then for any $n \geq 2$, take an element u_n of R such that $u_n \in X_n \setminus X_{n-1}$. Clearly, there is an infinite ascending chain $u_1 < u_1 + u_2 < u_1 + u_2 + u_3 < \cdots$, and hence ACC does not hold in A.

(2) If DCC does not hold for hereditary ideals of A, then there exists a strict descending chain of hereditary ideals of A, say, $Y_1 \supset Y_2 \supset \cdots$. Then for any $n \geq 1$, take an v_n such that $v_n \in Y_n \setminus Y_{n+1}$. Clearly, there is an infinite descending chain of elements of R, namely $\sum_{i\geq 1} v_i > \sum_{i\geq 2} v_i > \cdots$, where $\sum_{i\geq m} v_i \in Y_m$. Then DCC does not hold for elements of A. This completes proof. ■

Corollary 2.14. (1) *If ACC (DCC, respectively) holds for ideals of a po-semiring A, then ACC (DCC, respectively) also holds for elements of A.*

(2) *If ACC holds for elements of A, then ACC holds for principal annihilating ideals of A.*

(3) *For a po-semiring A, assume that there is the concept of an infinite sum of elements in A and any hereditary ideal I is closed under taking infinite sums. If further DCC holds for elements of A, then DCC holds for principal annihilating ideals of A.*

Corollary 2.15. *Let R be a commutative ring and denote $A = \mathbb{I}(R)$. Then ACC (DCC, respectively) holds for elements of A if and only if ACC (DCC, respectively) holds for hereditary ideals of A.*

The proof of the following result is routine and is omitted here.

Proposition 2.16 *For a po-semiring A and an element p of A, p is a prime element of A if and only if $< p >$ is a prime ideal of A.*

3. The graph $\Gamma(A)$ for a po-semiring A

For any multiplicative semigroup S with zero element 0, there is a zero divisor graph $\Gamma(S)$ whose vertices are nonzero divisors and there is an edge $x - y$ if $x \neq y$, $xy = 0$. Some fundamental properties of the semigroup graphs were given in [8]. For a semiring A, denote by $Z(A)$ the set of all nonzero multiplicative zero-divisors. The zero divisor graph of $(A, \cdot, 1)$ is also denoted as $\Gamma(A)$. Note that in a semiring, there exists two semigroup structures, and all known results on the zero divisor graphs of semigroups certainly holds for $\Gamma(A)$ (see [8]). Following [5], denote $\Gamma(\mathbb{I}(R)) = \mathbb{AG}(R)$ and call it *the annihilating* *ideal graph of R.* The graph was studied recently by quite a few authors, see[5, 2, 3].

Recall that a graph is called *complete* (*or discrete*) if every pair (or no pair) of vertices are adjacent. We denote a complete graph by K, a complete (or discrete) graph with n vertices by K_n (or D_n). An induced subgraph K of a graph G is called a *clique* if any two distinct vertices of K are adjacent and the *clique number* $\omega(G)$ of G is the least upper bound of the size of the cliques. Similarly, we denote by $K_{m,n}$ the complete bipartite graph with two partitions of sizes m, n. A complete bipartite graph $K_{1,n}$ is also called a *star graph.* Recall that a *cycle* in a graph is a path $v_1 - v_2 - \cdots - v_n$ together with an additional edge $v_n - v_1$ ($n \geq 3$). For any vertex x of G, let $N(x)$ be the vertices adjacent to x. If $|N(x)| = 1$, then x is called an *end vertex.* Recall a convenient construction from graph theory, *the sequential sum*

$$G_1 + G_2 + \cdots + G_r$$

of a sequence of graphs $G_1, G_2, \ldots, G_r$. The resulting graph $G_1 + G_2 + \cdots + G_r$ is obtained by adding an edge between each vertex of G_i and every vertex of G_{i+1} for all $i = 1, 2, \cdots, r-1$. We illustrate the construction in Figure 1 for the sequence of graphs D_2, K_1, K_1, D_3, where D_j is the discrete graph of j vertices. Such a resulting graph is a special case of two-star graphs which has two vertices as subcenters. Recall that a finite or an infinite graph G is called a *two-star graph* if $G \cong D_r + K_1 + K_1 + D_s$ ($\infty \geq r \geq 1 \leq s \leq \infty$).



Figure 1. Sequential sum of graphs

By [1, Theorem 2.2], for a commutative ring R, the zero divisor graph $\Gamma(R)$ is finite if and only if either R is a finite ring or an integral domain. The analog of the above result does not hold for a semigroup. However, such a result does exist for a class of po-semirings.

Theorem 3.1. *For a po-semiring A which is not integral, assume that a condition (1) or (2) as in Theorem 2.2 holds. Then $\Gamma(A)$ is a finite graph if and only if the po-semiring A is finite. In this case, $|V(\Gamma(A))| = |A| - 2$.*

Proof. If A is finite or integral, then $0 \leq |V(\Gamma(A))| < \infty$. The converse follows from Theorem 2.2. By Theorem 2.2, we also have $|V(\Gamma(A))| = |A| - 2$ in the case. ■

The following subtle example shows that condition (C_2) is crucial to Theorem 3.1. The construction in this example could be easily extended to obtain a finite or an infinite po-semiring A such that $\Gamma(A) \cong K_n$ for each natural number n. The example is also related to Theorem 2.2.

Example 3.2. *There exists an infinite po-semiring A such that the graph $\Gamma(A)$ is an isolated vertex.*

Let $A = \{0, 1, a, b_1, b_2, \cdots\}$ be a countable set with $|A| \geq 4$. Define the partial order and the addition as in Example 2.6, i.e., $0 < a < b_1 < b_2 < \cdots < 1$, $x + y = \max\{x, y\}$, $\forall x, y \in A$. Define a new multiplication by

$$0z = 0,\ 1z = z,\ (\forall z \in A),\ a^2 = 0,\ xy = \min\{\text{x,y}\} \text{ for other nonzero } x, y.$$

Then it is routine to check that A is a po-semiring. Clearly, $Z(A) = \{a\}$ but $|A|$ can be any cardinal number from 4 to $\aleph_0$. Note that DCC holds for elements of A, and that there exists no infinite set of orthogonal idempotents in A. Note also that condition (C_2) does not hold in A, but condition (C_3) does. Clearly $\Gamma(A)$ is a finite graph. Note that $\Gamma(A)$ is an infinite star graph for another po-semiring defined on the set A, which is constructed in Example 2.6.

Lemma 3.3. *Let $A = A_1 \times A_2$ be a direct sum of two semirings and denote $G = \Gamma(A)$. Then*

(1) *G contains no triangle if and only if one A_i is integral while $0 \leq |Z(A_j)| \leq 1$ for the other A_j.*

(2) *G contains no cycle if and only if one A_i is $\{0, 1\}$ while $0 \leq |Z(A_j)| \leq 1$ for the other A_j.*

(3) *G contains no quadrilateral if and only if one A_i is $\{0, 1\}$ while for the other A_j, either $0 \leq |Z(A_j)| \leq 1$, or $|Z(A_j)| = 2$ and there exists no nilpotent element in A_j.*

Proof. (1) $\Longleftarrow$: If both A_i are integral, then G is a complete bipartite graph and hence contains no triangle. In this case, one part contains $|A_1| - 1$ vertices and the other $|A_2| - 1$ vertices. If A_1 is integral and $Z(A_2) = \{x\}$, then $x^2 = 0$ and G is also a complete bipartite graph together with $|A_1| - 1$ end vertices adjacent to $(0, x)$. G clearly contains no triangles and is a bipartite graph, one part of which contains $2(|A_1| - 1)$ vertices and the other part $|A_2| - 1$ vertices.

$\Longrightarrow$: If neither A_1 nor A_2 is integral, assume $a_ib_i = 0$ with $a_i, b_i \in A_i \setminus \{0\}$ for $i = 1, 2$. Then in G there is a triangle $(0, a_2) - (a_1, 0) - (b_1, b_2) - (0, a_2)$. Now assume that A_1 is integral. If $|Z(A_2)| \geq 2$, then there exists in G a triangle $(1, 0) - (0, a) - (0, b) - (1, 0)$.

(2) $\Longleftarrow$: This follows from the sufficiency proof of (1).

$\Longrightarrow$: If $|A_i| \geq 3$ for all i, then there is in G a square $(1, 0) - (0, a) - (b, 0) - (0, 1) - (1, 0)$. Assume $A_1 = \{0, 1\}$. If $|Z(A_2)| \geq 2$, then there exists in G a triangle $(1, 0) - (0, a) - (0, b) - (1, 0)$.

(3) $\Longleftarrow$: Assume $A_1 = \{0, 1\}$. If $0 \leq |Z(A_j)| \leq 1$, then G clearly contains no cycle. If $|Z(A_j)| = 2$ and there exists no nilpotent element in A_j, then there exists a triangle in G

but there is no square in G.

$\Longrightarrow$: If $|A_i| \geq 3$ for all i, then there is in G a square. Now assume $A_1 = \{0,1\}$. If $|Z(A_2)| \geq 3$, then there is a path $u - v - w$ in $\Gamma(A_2)$. Then there is in G a square $(1,0) - (0,u) - (0,v) - (0,w) - (1.0)$. Now assume $|Z(A_2)| = 2$. If there is a nilpotent element in A_2, assume $u^2 = 0$. Then there is a square $(1,0) - (0,u) - (1,u) - (0,v) - (1,0)$. This completes the verification. ■

Minimal elements of a po-semiring A play a special role in the graph $\Gamma(A)$. More precisely, we have the following.

Lemma 3.4. *Let A be a po-semiring and denote $G = \Gamma(A)$. Assume further $Z(A) \neq \emptyset$. Then*

(1) *Let $a - u - b$ be a path in G. If it is contained in neither triangle nor quadrilateral, then u is a minimal element of A.*

(2) *Each minimal element is a zero divisor of A, and thus the clique number $\omega(G)$ is greater than or equal to the number of minimal elements of A.*

(3) *Let u be a minimal element of A. Then*

(3.1) $d(u,x) \leq 2$, $\forall x \in V(G)$.

(3.2) *For each clique K of G, either $V(K) \subseteq N(u)$ or $V(K) \setminus \{v\} \subseteq N(u)$ for a vertex v in $V(K)$.*

Proof. (1) Assume to the contrary that u is not a minimal element of A. Assume further $0 < v < u$. If $v = a$, then there is a triangle $a - u - b - a$. If $v \neq a, b$, then there is a square $a - u - b - v - a$.

(2) Let $x \in Z(A)$. First, for any minimal element e of A, either $ex = 0$ or $ex = e$. In particular, if $ex \neq 0$, then $N(x) \subseteq N(e)$ in $\Gamma(A)$. Thus each minimal element of A is a zero divisor of A. It implies that the clique number of $\Gamma(A)$ is greater than or equal to the number of minimal elements of A.

(3) By (2), $u \in V(G)$. $\forall u \neq x \in V(G)$, if $ux \neq 0$, then $N(x) \subseteq N(u)$. This shows $d(u,y) \leq 2$, $\forall y \in V(G)$. (3.2) then also follows from the observation. ■

We remark that for the po-semiring $A = \{0,1\}^{(n)}$ (see Example 3.6 for the definition), the clique number $\omega(\Gamma(A))$ is identical with the number of minimal elements of A.

Here is the main result of this section.

Theorem 3.5. *Let A be either a commutative ring or a po-semiring satisfying condition (C_3). Let $\Gamma(A) = G$.*

(1) *If G contains no cycle, then G is either a star graph or a two-star graph $K_1 + K_1 + K_1 + D_r$.*

(2) $\Gamma(A) \cong K_1 + K_1 + K_1 + D_r$ *if and only if $A \cong \{0,1\} \times S$, where S is a po-semiring*

with $|Z(S)| = 1$*. In this case,* $r = |S| - 2$*.*

Proof. (1) By [8, Proposition 1.3], for any commutative semigroup S with 0, if $\Gamma(S)$ is a tree, then $\Gamma(S)$ is a connected subgraph of a two-star graph. It holds in particular for po-semirings. If further the diameter of G is two, then G is clearly a star graph. Now assume $diam(G) = 3$ and assume $x - u - v - y$ be a path in G. Then G is a two-star graph with subcenters u, v. By [12, Corollary 2.13], we can assume $u^2 = u$. Then $v^2 = 0$.

If A is a ring, then there is a ring isomorphism $A \cong Au \times A(1-u)$. The result follows from Lemma 3.3(2).

If A is a po-semiring, then by Lemma 3.4(1), both u and v are minimal element of A. Since condition (C_3) holds, there exists an idempotent w such that $A = Au \times Aw$, where $w + u = 1, wu = 0$. By Lemma 3.3(2), we can assume $Au = \{0, u\}$ and $0 \leq |Z(Aw)| \leq 1$. By the sufficiency proof of Lemma 3.3(1), we have $|Z(Aw)| = 1$ and $G \cong K_1+K_1+K_1+D_r$, where $r = |Aw| - 2$.

(2) This follows from the proof of (1) and the proof of Lemma 3.3(2). ■

The following example shows that Theorem 3.5 is the best possible result for po-semirings satisfying condition (C_3).

Example 3.6. *For any* $1 \leq r \leq \aleph_0$*, there exists a po-semiring* A *such that condition* (C_3) *holds and* $\Gamma(A) \cong K_1 + K_1 + K_1 + D_r$*.*

In fact, for any po-semirings A_i $(1 \leq i \leq n)$, define a partial order $\leq$ in the semiring $A_1 \times A_2 \times \cdots \times A_n$ in a natural way, i.e., $(x_1, \ldots, x_n) \leq (y_1, \ldots, y_n)$ if $x_i \leq y_i$ for all i. Then $A_1 \times A_2 \times \cdots \times A_n$ is a po-semiring, called the *direct product of* $A_1, \cdots, A_2$. When $A_1 = \cdots = A_n = A$, denote by $A^{(n)}$ the above direct product. Take $A_1 = \{0, 1\}$ and

$$A_2 = \{0, 1, a, b_1, b_2, \cdots\}$$

as in Example 3.2. Then $\Gamma(A_1 \times A_2) = K_1 + K_1 + K_1 + D_r$, where $r = |A_2| - 2$ can be any cardinal number from 1 to $\aleph_0$. Clearly, condition (C_3) holds in $A_1 \times A_2$ since there are only two minimal elements (i.e., $(1, 0), (0, a)$).

Example 3.7. $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_4) = \Gamma(\mathbb{Z}_2[x]/(x^2)) = K_1 + K_1 + K_1 + D_2$. For any commutative ring R, by [2, Theorem 2], if $\mathbb{AG}(R)$ is a tree, then it is either a star graph or the two-star graph $K_1 + K_1 + K_1 + K_1$.

Corollary 3.8. *If condition* (C_1) *holds in a po-semiring* A *and there exists no infinite set of orthogonal idempotents in* A*, then*

(1) $\Gamma(A)$ *is either a star graph or the two-star graph* $K_1 + K_1 + K_1 + K_1$*, if further the graph* $\Gamma(A)$ *contains no cycle.*

(2) $\Gamma(A) \cong K_1+K_1+K_1+K_1$ *if and only if* $A \cong \{0,1\} \times S$*, where* S *is the po-semiring*

$\{0, a, 1\}$ with $a^2 = 0$.

Remark 3.9. (1) It is easy to see that there are isomorphically two semirings A with $A = \{0, 1\}$, namely the ring $\mathbb{Z}_2$ and the po-semiring $\mathbb{I}(K)$ for any field K. Also, there exist exactly two po-semirings A with $|A| = 3$, namely $A = \{0, a, 1\}$ with partial order $0 < a < 1$, and with multiplication defined by either $a^2 = 0$ or $a^2 = a$ respectively.

(2) In a semiring A, if $|Z(A)| = 1$, then either the ring $\mathbb{Z}_2[x]/(x^2)$ or $\mathbb{Z}_4$ can be embedded into A. While in the ring $\mathbb{Z}_n$, if there is a semiring S contained in $\mathbb{Z}_n$ such that $|Z(S)| = 1$, then $n = 4$ and $S = \mathbb{Z}_4$.

(3) It is natural to ask the following question: Can any two-star graph be realized as the zero divisor graph of a po-semiring when condition (C_3) is dropped? The answer is yes and see [14] for a complete answer.

4. The structure of a po-semiring A with small $|Z(A)|$

In this section, motivated by Lemma 3.3 and Theorem 3.5, we study the structure of a po-semiring A with small $Z(A)$. We first have the following Lemma.

Lemma 4.1. *Let A be a po-semiring.*

(1) If $Z(A) = \{c\}$, then $c^2 = 0$, c is the least nonzero element of A and is a prime element of A.

(2) If $Z(A) = \{c, u\}$, then exactly one of the following holds:

(2.1) Both c and u are minimal elements of A, $c < x$, $u < x$, $\forall x \in A \setminus \{0, c, u\}$, $u^2 = u$ and $c^2 = c$. Furthermore, both c and u are prime elements of A.

(2.2) c is the least nonzero element of A, and $c^2 = 0$. u is a prime element of A, and $u < p$ for each prime element $p \neq c, u$.

Proof. (1) If $Z(A) = \{c\}$, then clearly $c^2 = 0$. Now if $xy \leq c$ for some $x, y \in A$, then $x(xy^2) = 0$. If further $x \not\leq c$, then $xy^2 = 0$ and hence $y^2 = 0$. Then $y \leq c$. This shows that c is a prime element of A. Now for any $x \in A \setminus \{0, c\}$, clearly $c = xc$ and hence $c < x$. Thus c is the least nonzero element of A.

(2) Let $Z(A) = \{c, u\}$. First, assume that c and u are incomparable. Then $c + u \notin \{c, u\}$ and hence $c^2 = c, u^2 = u$. It follows that $xc = c$, $xu = u$, $\forall x \neq 0, c, u$, and hence $c < x, u < x$ hold for each $x \neq 0, c, u$. It implies that both c and u are minimal elements of A. To verify that c is a prime element, assume $xy \leq c$ and $y \not\leq c$. If $y = u$, then $x \leq c$. If further $y \neq u$, then $0 = (xy)u = xu$ and so $x \leq c$. Thus c is a prime element of A. By symmetry, u is also a prime element of A.

Next we assume $c < u$. Then $c^2 = 0$. For any $x \in A \setminus \{0, c, u\}$, clearly $c = xc$ and hence $c < x$. Thus c is the least nonzero element of A. For any $x \not\leq u, y \not\leq u$, $xy \not\leq u$ since otherwise, $0 = (xy)c = x(yc) = c$, a contradiction. Thus u is a prime element of A. Finally, for any prime element $p \neq c, u$, if $u^2 \in \{0, c\}$, then $u^3 = 0$ and hence $u < p$. If

$u^2 = u$, then $pu \neq c$ and hence $u = pu < p$. ■

Theorem 4.2. (1) *A is a po-semiring with $|Z(A)| = 1$ if and only if there exist an integral po-semiring A_1 and an element $c \notin A_1$ such that $A \cong \{c\} \cup A_1$, where the partial order of $\{c\} \cup A_1$ is extended from A_1 by adding $0 < c < a$ ($\forall a \in A_1^*$), and the commutative addition and multiplication in $\{c\} \cup A_1$ are extended respectively from that of the po-semiring A_1, and the following conditions are fulfilled:*

$$0 + c = c,\ c + c = c,\ c^2 = 0,\ c + y = y,\ cy = c\,(\forall y \in A_1^*).$$

(2) *A is a po-semiring such that $|Z(A)| = 2$ and $Z(A)^2 \neq 0$ if and only if one of the following holds:*

(2.1) *There exist an integral po-semiring A_1 and elements $c, u \notin A_1$ such that $A \cong \{c, u\} \cup A_1$, where A_1 has a least nonzero element a_0, the partial order of $\{c, u\} \cup A_1$ is extended from A_1 by adding $0 < c < x$ and $0 < u < x$ ($\forall x \in A_1^*$), and the commutative addition and multiplication in $\{c\} \cup A_1$ are extended respectively from that of the po-semiring A_1, and the following conditions are fulfilled:*

$$c + u = a_0,\ 0 + x = x,\ x + x = x,\ x + y = y\,(\forall x \in \{c, u\},\ \forall y \in A_1^*),$$

$$u^2 = u,\ c^2 = c,\ cu = 0 = 0x,\ xy = x\,(\forall x \in \{c, u\},\ \forall y \in A_1^*).$$

(2.2.) *There exist an integral po-semiring A_1 and elements $c, u \notin A_1$ such that $A \cong \{c, u\} \cup A_1$, where the partial order of $\{c, u\} \cup A_1$ is extended from A_1 by adding $0 < c < u < a$ ($\forall a \in A_1^*$), and the commutative addition and multiplication in $\{c\} \cup A_1$ are extended respectively from that of the po-semiring A_1, and the following conditions are fulfilled:*

$$x + y = max\{x, y\},\ \forall x \in \{c, u\},\ \forall y \in \{c, u\} \cup A_1,$$

$$c^2 = 0,\ cu = 0 = 0x,\ u^2 \in \{c, u\},\ xy = x\,(\forall x \in \{c, u\},\ \forall y \in A_1^*).$$

Proof. (1) $\Longrightarrow$: Let A be a po-semiring and denote $A_1 = A \setminus Z(A)$. Then A_1 is an integral sub-po-semiring of A, and the rest results follow from Lemma 4.1(1).

$\Longleftarrow$: It is routine to check that $\{c\} \cup A_1$ is a po-semiring under the assumptions. Clearly $|Z(A)| = 1$.

(2)$\Longrightarrow$: Assume that A is a po-semiring such that $|Z(A)| = 2$. Assume further $Z(A) = \{c, u\}$, and set $A_1 = A \setminus \{c, u\}$.

(2.1) If c and u are incomparable, then $c + u \in A_1^*$ and $Z(A)^2 \neq 0$ by Lemma 4.1(2). Assume $c + u = a_0$. By the proof of Lemma 4.1(2), we have $a_0 y = (c + u)y = c + u = a_0, \forall y \in A_1^*$. Thus a_0 is the least nonzero element of A_1. Since both c and u are prime elements of A by Lemma 4.1(2), A_1 is an integral sub-po-semiring of A. The other statements also follow directly from Lemma 4.1(2).

(2.2) In the following, assume that c and u are comparable, assume further $c < u$. Then $c^2 = 0$, and c is the least nonzero element of A. Thus $ac = c$ for any nonzero element a in A_1^*. If $Z(A)^2 \neq 0$, then $u^2 \neq 0$. This implies $au = u, \forall a \in A_1^*$. Then $u < a$, $\forall a \in A_1^*$. Since u is a prime element of A, $u \notin A_1^* A_1^*$. Clearly, $c \notin A_1^* A_1^*$. Thus A_1 is an integral sub-po-semiring of A. This completes the necessary part of the proof.

$\Longleftarrow$: It is not hard to check that $\{c, u\} \cup A_1$ is a po-semiring under either assumption. Clearly, $|Z(A)| = 2$ and $Z(A)^2 \neq 0$ hold in either case. Note that (C_3) holds in Case (2.2). ■

We remark that Theorem 4.2 (2.1) gives a complete characterization of po-semirings A with $|Z(A)| = 2$, in which there exist no nilpotent elements. In particular, we have the following.

Corollary 4.3 (1) *A po-semiring A satisfies condition (C_3), $|Z(A)| = 2$ and $Z(A)^2 \neq 0$ if and only if either $A \cong \{0,1\} \times \{0,1\}$, or A is the po-semiring constructed in Theorem 4.2(2.2).*

(2)*A po-semiring A satisfies condition (C_3), $|Z(A)| = 2$ and there exist no nilpotent elements in A if and only if $A \cong \{0,1\} \times \{0,1\}$.*

Note that the po-semiring $\mathbb{I}(R)$ always satisfies condition (C_3). Applying Corollary 4.3 to the po-semiring $\mathbb{I}(R)$, we have the following corollary which is essentially the same with [3, Theorem 3] by [11, Theorem 2.1].

Corollary 4.4. *For any commutative ring R, The following statements are equivalent:*

(1) *Either $R \cong F_1 \times F_2$ for some fields F_i, or R is local with two nontrivial ideals.*

(2) *The annihilating graph $\mathbb{AG}(R)$ is isomorphic to the complete graph K_2.*

(3) *Either $R \cong F_1 \times F_2$ for some fields F_i, or R is a local ring with the maximal ideal $J(R) = R\alpha$ for some α satisfying $\alpha^3 = 0$ and $\alpha^2 \neq 0$.*

Proof. (1) $\Longrightarrow$ (2) : Clear.

(2) $\Longrightarrow$ (3) : Assume $\mathbb{AG}(R) \cong K_2$. Then R has exactly two proper ideals by [5, Theorem 1.4]. In particular, R is artinian and thus the ideal $J(R)$ is finitely generated. It follows that $J(R) = R\alpha$ for some $\alpha \in J(R)$.

If $J(R)^2 = 0$, then either $R \cong F_1 \times F_2$ for some fields F_i, or R is a local ring by Lemma 4.1(2) and Corollary 4.3. If further R is a local ring, then $J(R) = U(R)\alpha$ since $J(R)\alpha = 0$. Under the assumption, R has exactly one nontrivial ideal, a contradiction. The contradiction shows $R \cong F_1 \times F_2$ if $J(R)^2 = 0$.

Now assume $J(R)^2 \neq 0$. By Corollary 4.3, R is a local ring with exactly two nontrivial ideals. Since $J(R)^2 \neq J(R)$ by Nakayama Lemma, $J(R)$ and $J(R)^2$ are the nontrivial ideals of R by Theorem 4.2(2.2). Hence $J(R)^3 = 0$ and $J(R)^2 \neq 0$. In the case, R is a local ring whose unique maximal ideal is $J(R) = R\alpha$, where $\alpha^3 = 0, \alpha^2 \neq 0$.

(3) $\Longrightarrow$ (1) : Now assume that R is a local ring with $J(R) = R\alpha$, where $\alpha^3 = 0, \alpha^2 \neq 0$. Clearly, $R\alpha \neq R\alpha^2$ and $R\alpha \cdot R\alpha^2 = 0$. $R = U(R) \cup R\alpha$ and hence $R\alpha = U(R)\alpha \cup U(R)\alpha^2 \cup \{0\}$. Thus for any $\beta \in J(R)$, $R\beta = R\alpha$ if $\beta \in U(R)\alpha$, while $R\beta = R\alpha^2$ if $0 \neq \beta \in U(R)\alpha^2$. It shows that $R\alpha$ and $R\alpha^2$ are all the possible nontrivial ideals of R. ∎

The complete isomorphic classification of finite local rings with at most three nontrivial ideals will be discussed in a separate paper, see [13].

Note that a po-semiring A with $|Z(A)| = 1$ always satisfies condition (C_3). By Theorem 4.2, *the structure of a po-semiring A is completely determined by structures of integral po-semirings if A satisfies one of the following conditions:*

(1) $|Z(A)| = 1$.

(2) $|Z(A)| = 2$ and $Z(A)^2 \neq 0$.

On the other hand, the structure of A with $Z(A)^2 = 0$ seems to be a little bit more complicated. Let A be a po-semiring with $Z(A) = \{c, u\}$ and $Z(A)^2 = 0$. Set $A_1 = A \setminus Z(A)$. Then A_1 is an integral sub-po-semiring (see Proposition 4.5 below). Clearly, $u < x + u < 1$ for any $x \notin \{0, 1, c, u\}$. By Lemma 4.1(2.2), c is the least nonzero element of A, so $cA_1^* = \{c\}, c + x = x, \forall x \in A^*$. But it is hard to determine the partial order between u and elements of A_1^*. By Examples 4.6 and 4.7 below, it seems that Lemma 4.1(2.2) is the best possible result.

Proposition 4.5. *A is a po-semiring with condition (C_3), $|Z(A)| = 2$ and $Z(A)^2 = 0$ if and only if there exist an integral po-semiring A_1 and elements $c, u \notin A_1$, such that $A \cong Z(A) \cup A_1$, where $Z(A) = \{c, u\}$, the addition and multiplication in $Z(A) \cup A_1$ are extended respectively from that of the po-semiring A_1, and the following conditions are fulfilled:*

(1) $0 + x = x \, (\forall x \in Z(A)), c + y = y \, (\forall y \in A^*), u + 1 = 1, u + u = u,$ *and* $u + (x + y) = (u + x) + y$, $u + (u + x) = u + x \, (\forall x, y \in A_1^*)$.

(3) $Z(A)^2 = 0, 0Z(A) = 0, cx = c \, (\forall x \in A_1^*), 0 \notin u \cdot A_1^*$.

(4) $xu = u$ *and* $yu = u$ *implies* $(xy)u = u$.

(5) $x \geq y$ *in* A_1 *and* $yu = u$ *implies* $xu = u$.

(6) *For any* $y, z \in A_1$, $x(y + u) = xy + xu$. *Also,* $uy = c = uz$ *implies* $u(y + z) = c$.

Proof. $\Longrightarrow$: By Lemma 4.1, it suffices to show that $A_1 = A \setminus Z(A)$ is a sub-po-semiring. In fact, if $xy \in \{c, u\}$ for some $x, y \in A_1^*$, then we have $xc = x(yc) = (xy)c = 0$, giving a contradiction.

$\Longleftarrow$: We omit the detailed verification here. ∎

Next we use Proposition 4.5 to construct some po-semirings A with $|Z(A)| = 2$, $Z(A)^2 = 0$.

Example 4.6. Let $A = \{0, 1, c, u, b_1, b_2, \cdots\}$ be a poset with $4 \leq |A| \leq \aleph_0$, where

$0 < c < u < b_1 < b_2 < \cdots$. Define the binary addition by max operation. Define three commutative multiplications by

$$0 \cdot x = 0,\ 1 \cdot x = x\,(\forall x \in A),\ cu = 0,\ u^2 \in \{0, c, u\},\ y \cdot z = min\{y, z\}(\text{for other } y, z).$$

Then it is easy to check that A is a po-semiring for each multiplication. Clearly, $|Z(A)| = 2, Z(A)^2 = 0$.

Example 4.7. Let $Z(A) = \{c, u\}$ and $A = Z(A) \cup A_1$, where $A_1 = \{0, 1, b_1, b_2, \cdots\}$ is a chain $0 < b_1 < b_2 < \cdots < 1$, with *max* as the binary addition and with multiplication $b_i b_j = b_1$ $(\forall i, j \geq 1)$. Then clearly A_1 is an integral po-semiring. Fix a positive integer $n > 1$ and extend the partial order of A_1 to A by $0 < c < u < b_n$. Extend the commutative addition to A by

$$0 + x = x,\ 1 + x = 1\,, x + x = x\,(\forall x \in \{c, u\}, c + y = y\,(\forall y \in A^*)$$

and

$$u + b_i = b_n\,(1 \leq i \leq n-1), u + b_j = b_j\,(j \geq n).$$

Extend the commutative multiplication to A by

$$0Z(A) = 0 = Z(A)^2,\ ub_i = c\,(\forall i).$$

Note that $c + y = y\,(\forall y \in A^*)$ implies that c is the least element of A, thus $c^2 = 0$ means that condition (C_3) holds. Note that $0 + x = x$, $x + 1 = 1$ implies $0 \leq x \leq 1$. Then it is easy to apply Proposition 4.5 check that A is a po-semiring with $|Z(A)| = 2, Z(A)^2 = 0$. *Note that u and b_i are incomparable for any i with $1 \leq i \leq n-1$, while by Lemma 4.1(2), $u < p$ for any prime element p with $p \neq c, u$.*

For any po-semiring A, recall that $A^{(r)}$ is the direct product of r copies of A. We have the following.

Proposition 4.8. *A is a po-semiring, in which both DCC and condition (C_3) holds for elements of A, if and only if there exist an integer $n \geq 1$ and a po-semiring A_1 such that either $A \cong A_1$ or $A \cong \{0, 1\}^{(n)} \times A_1$, where $c^2 = 0$ for each minimal element c of A_1 and DCC holds for elements of A_1.*

Proof. $\Longleftarrow$: Let A_1 be a po-semiring, and assume that DCC holds for elements of A_1 and $c^2 = 0$ for each minimal element c of A_1. Then there exists at leat one minimal element c in A and hence $Z(A_1) \neq \emptyset$. If $A \cong A_1$, then clearly condition (C_3) holds in A. If $A \cong \{0, 1\}^n \times A_1$ for some finite $n \geq 1$, then both DCC and condition (C_3) also hold for elements of A.

$\Longrightarrow$: Assume that DCC holds for elements of A. Then there exists at leat one minimal element in A. If further $c^2 = 0$ for each minimal element c of A, then $Z(A) \neq \emptyset$ and condition (C_3) holds in A. Then we take $A_1 = A$. In the following assume that there exists an idempotent minimal element e_1 in A. Then by condition (C_3), there exists a nonzero idempotent $f_1 \in A$ such that $A \cong \{0,1\} \times Af_1$. Clearly, both DCC and condition (C_3) hold for elements of the po-semiring Af_1 and an induction shows that $A \cong \{0,1\}^{(n)} \times A_1$ for some $n \geq 1$, where A_1 is a po-semiring in which $c^2 = 0$ for each minimal element c of A_1 and DCC holds for elements of A_1. $\blacksquare$

References

[1] D.F. Anderson and P.S. Livingston. The zero-divisor graph of a commutative ring, **J. Algebra 217**(1999) 434 – 447.

[2] G. Aalipour, S. Akbari, R. Nikandish, M.J. Nikmehr and F. Shaveisi. Minimal prime ideals and cycles in annihilating-ideal graphs, **Rocky Mountain J. Math.** (accepted).

[3] G. Aalipour, S. Akbari, M. Behboodib, R. Nikandish M.J. Nikmehr and F. Shaveisi. The Classification of the Annihilating-Ideal Graph of a Commutative Ring. **Algebra Colloquium** (accepted)

[4] M.F. Atiyah and I.G. MacDonald. *Introduction to Commutative Algebra*, **Addison-Wesley**, Reading, MA, 1969.

[5] M. Behboodi and Z. Rakeei. The annihilating-ideal graph of commutative rings I, **Journal of Algebra and its Applications** (accepted).

[6] S. Bistarelli, U. Montanari, and F. Rossi. Semiring-based constraint satisfaction and optimization, **Journal of the ACM**, **44:2**(1997) 201 – 236.

[7] S. Bistarelli, U. Montanari, F. Rossi, Schiex, T.; G. Verfaillie, H. Fargier. Semiring-based CSPs and valued CSPs: frameworks, properties, and comparison, **Constraints 4:3** (1999) 199 – 240.

[8] F.R. DeMeyer, T. McKenzie, and K. Schneider. The zero-divisor graph of a commutative semigroup, **Semigroup Forum 65**(2002) 206 – 214.

[9] T.Y. Lam. A First Course in Non-Commutative Rings, **Springer-Verlag**, New York, Inc 1991.

[10] D.C. Lu and T.S. Wu. On the normal ideals of exchange rings, **Siberian Math.J. 49**(2008) 663 – 668.

[11] K.R., McLean. Commutative artinian principal ideal rings. **Proc. London Math. Soc.** (3) **26** (1973), 249 – 272.

[12] T.S. Wu and D.C. Lu. Sub-semigroups determined by the zero-divisor graph, **Discrete Math. 308**(2008) $5122-5135$.

[13] T.S. Wu and D.C. Lu. Finite local rings with at most three nontrivial ideals. Preprint.

[14] H.Y. Yu and T.S. Wu. On realizing zero-divisor graphs of po-semirings. Preprint.